

Instructions For Use MTF-IFU

Rev. Date: Dec. 21, 2013

Revision: 4

Page 1 of 1

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

Michel's Transport Fluid

Description: Michel's Transport Fluid is intended for use as a transport media of specimens (such as renal biopsies

and lymph nodes) for immunofluorescence studies. Samples may remain in media for up to 5 days at room temperature. This reagent is not a fixative and is not suitable for transporting live cells for flow cytometry. Prior to processing, specimens should be washed in 3 changes of Michel's Transport Wash

Buffer (cat.# MTW) for 10 minutes each.

Uses/Limitations: Not to be taken internally.

For In-Vitro Diagnostic use only.

Histological applications.

Do not use if reagents become cloudy. Do not use past expiration date.
Use caution when handling reagents.

Non-Sterile.





Availability: <u>Item #</u> <u>Volume</u>

MTF125 125 ml MTF500 500 ml MTF999 1000 ml MTF-10000 10 Liters MTF-20000 20 Liters

Storage: Store at 2-8 ℃.

Precautions: Avoid contact with skin and eyes.

Harmful if swallowed.

Follow all Federal, State, and local regulations regarding disposal.

Procedure:

- 1. Pour reagent into appropriate sized container.
- 2. Place sample in container.
- 3. Add enough additional Transport Fluid (if required) to completely cover specimen.
- 4. Transport sample to evaluation site immediately (packing in ice is not required).

References:

1. Vaughn Jones, S.A., et al. The use of Michel's transport medium for immunofluorescence and immunoelectron microscopy in autoimmune bullous diseases. Journal of Cutaneous Pathology, 1995 August; 22(4): pages 365-370.

Storage: 2° C



8° C

ScyTek Laboratories, Inc. 205 South 600 West Logan, UT 84321 U.S.A.



EC REP EmergoEurope (31)(0) 70 345-8570 Molsnstraat 15 2513 BH Hague, The Netherlands

Doc: IFU-Template2-8rev2