

Instructions For Use		
TBD-IFU		
Rev. Date: Jan. 19, 2011	Revision: 3	Page 1 of 1

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

Tris Buffered Saline (10X) pH 7.5

Description: Tris Buffered Saline (10X) pH 7.5 has been formulated in an easy-to-use 10X concentration for use with

immunolabeling techniques as a wash buffer. Made using only ACS Reagent grade materials and water that has been both deionized and distilled provides assurance that optimal results can be expected. The need to weigh and dissolve individual components is eliminated and replaced with a simple dilution.

Contents: Tris buffered saline in deionized/distilled water. pH of diluted buffer is 7.5±0.2. Concentration at final

dilution is Tris: 0.065M, NaCI: 0.2M, KCI: 0.003M.

Uses/Limitations: Not to be taken internally.

For In-Vitro Diagnostic use only. Do not use if reagent becomes cloudy. Do not use past expiration date. Use caution when handling reagents.

Non-Sterile

Availability: <u>Item # Volume</u> <u>Diluted Volume</u>

 TBD500
 500 ml
 5 liters

 TBD999
 1000 ml
 10 liters

 TBD010
 10 liters
 100 liters

 TBD-20000
 20 liters
 200 liters

Storage: Store at 18-25 ℃.

Precautions: Avoid contact with skin and eyes.

Harmful if swallowed.

Follow all Federal, State, and local regulations regarding disposal.

Procedure:

- 1. Mix 1 part Tris Buffered Saline with 9 parts deionized and/or distilled water.
- 2. For optimal results, prepare fresh buffer daily.





