

Anti-Human CD90 (Thy-1) / Fibroblast Marker Mouse Monoclonal Antibody - FITC-conjugated

Product Information

Catalog-No.:	DIA-120 (200µq)	Species Reactivity	
Tests:	Minimum 100 Tests (10 µl/ Test)	Human	
Clone: Isotype: Specificity:	AS02 Mouse IgG1, kappa Human CD90 (Thy-1) (nerve cells, few CD34 ⁺ blood stem cells, fibroblasts)	No reaction with Rat (cultured fibroblasts, FC), Mouse (fibrosarcom cell line L929, FC), Rabbit (skin, IH), Pig (skin, IH), Monkey (skin, IH) Others not tested	
Immunogen:	Human skin fibroblasts	Applications	Dilution
Physical State:	Lyophilized powder	Flow Cytometry	1:50 – 1:500
Reconstitution: Presentation:	After opening, restore to 1 ml with sterile distilled water. FITC-conjugated Mouse IgG1 in PBS with 2% BSA, 0.05% NaN ₃ , pH 7.4. The anti- body was purified from culture supernatant by mouse IgG-specific affinity chromatog- raphy. FITC was coupled using standard procedures and unbound FITC was elimi- nated by gel chromatography.		100 μ l of an appropriate dilution for a suspension of 5 x 10 ⁵ cells
		(Immunofluorescence)	Use of unconjugated antibody with secondary detection antibody recommended.
		The indicated dilutions are general recommendations. For special applications optimal working dilutions should be de- termined individually.	

Reactivity

The monoclonal antibody clone AS02 reacts specifically with human CD90 (Thy-1), a GPI-anchored glycoprotein of the immunoglobulin superfamily with a molecular weight of 25 – 35 kDa (3). CD90 (Thy-1) in human is primarily expressed by nerve cells, additionally in a sub-population (20%) of CD34⁺ blood stem cells and in various fibroblasts (1-8). In contrast to mouse and rat it is not expressed by thymocytes and peripheral blood T cells. AS02 recognizes fibroblasts (1-8). In contrast to mouse and rat it is not expressed by thymocytes, resting micro-/macrovascular endothelial cells and components of the extracellular matrix like fibronectin, collagen type I, III, IV and laminin. Thus, AS02 is especially suitable for the specific detection and cell separation of human fibroblasts (2). Activated endothelial cells, after stimulation in vitro or in tissue sections of inflamed tissue in vivo, bind AS02 with different intensities, implying that CD90 (Thy-1) can be considered to be an activation marker of human endothelial cells. Recent papers demonstrate the binding of AS02 with activated microvas cular endothelial cells (4, 5), with a subpopulation of lung fibroblasts (6), specialized lymph node fibroblasts, but not lymphocytes (7) and normal or inflamed synovial fibroblasts in human (8). In immunoblot analysis AS02 detects no nred uced CD90 at 30 kDa (1, 3).

Reactivity of AS02 in immunohistology

Skin	Fibroblasts	+	Lymph Nodes Thyroid Gland/ Liver/ Kidney/ Gall Bladder	Fibroblasts	+
	Keratinocytes	-		Lymphocytes	-
	Endothelial Cells	-		Fibroblasts	+
	Smooth Muscle Cells	-		Epithelial Cells	-
	Glandular Cells	-		Thyrocytes	-
	Macrophages	-		Liver Parenchyma	-
	Langerhans Cells -	Kidney Parenchyma	-		
Cartilage	Chondrocytes	-		Tubulus Epithelial Cells	(+)
			Placenta	Fibroblasts (Mesenchyma)	+
Muscles	Smooth Muscle Cells	-		Endothelial Cells	-
	Heart Muscle Cells	-		Epithelial Cells	-
	Sceletal Muscle Cells	-	Brain	Nerve Cells	+

Legend: + strong binding; (+) recognition of few cells; - no binding

For in vitro research use only. Not for diagnostic or therapeutic use.







Reactivity of AS02 in flow cytometry



Binding of the monoclonal antibody AS02 to Dermal Fibroblasts in flow cytometrie. AS02 does not stain Lymphocytes, Keratinocytes, Monocytes, Granulocytes, Thrombocytes, HUVEC and HDMC. AS02 (------) compared to a an appropriate cell marker (------); HUVEC = (Macrovascular) Human Umbilical Vein Endothelial Cells, HDMEC = Human Dermal Microvascular Endothelial Cells.



Flow cytometric analysis of the binding capacity of mAb AS02 with different cell types. Staining of different types of human cells with ASO2 (------); HUVEC = (Macrovascular) Human Umbilical Vein Endothelial Cells, HDMEC = Human Dermal Microvascular Endothelial Cells.

For in vitro research use only. Not for diagnostic or therapeutic use.



BIOZOL GmbH Leipziger Str.4 D-85386 Eching Germany

www.biozol.de info@biozol.de +49 (0)89 - 3799 666 6







Flow cytometric analysis of fibroblasts of different origin with mAb AS02.

Staining of different types of human fibroblasts with AS02 (-------); only slight differences in the average fluorescence in tense ness and therefore the binding affinity of monoclonal fibroblast antibody clone AS02 show the high affinity of this antibody to fibroblasts of different origin.

Instructions for Use

Applications

- Flow Cytometry
- Laser Scanning Cytometry

Immunofluorescence (ethanol, methanol or acetone fixed cells and sections of frozon tissue only: formal debyde fixed paraffin om

of frozen tissue only; formaldehyde-fixed paraffin embedded material is NOT stained by AS02)

1. Immunofluorescence

- 1. Cells: fix cells in ice-cold acetone for 5 minutes, allow to air dry. Frozen sections: warm slides at room temperature for 30 and fix in ice-cold acetone for 10 minutes, air dry for 5 (likewise ice-cold ethanol or methanol are suitable for fixation)
- 2. Rinse cells/sections in 3 changes of washing buffer; 5 minutes each
- 3. Optionally block nonspecific binding with 5% FCS for 20 at room temperature
- Add 50 100 μl of FITC-conjugated anti-CD90 antibody AS02 (DIA 120), diluted 1:50 to 1:100 in PBS/1% BSA
- 5. Incubate at room temperature for 60 minutes
- 6. Rinse slides in 3 changes of washing buffer; 5 minutes each
- 7. Counterstain slides with DAPI
- 8. Coverslip with mounting medium (e.g. 80-90% glycerol/PBS + 2,5% DABCO) and seal with nail polish
- 9. Store slides in dark at 4 °C

2. Flow Cytometry

- Detach cultured fibroblasts by 0.025% trypsin/0.01% EDTA for 5 10 minutes at 37°C or by using 2.5 mM/L EDTA/PBS
- 2. Wash 2 x with cold PBS; centrifugation at 200 x g for 5 minutes each
- Resuspend 2 5 x 10⁵ cells in 100 µl of a 1:50 1:500 dilution of anti-CD90 antibody AS02 (DIA 120) in PBS/1% BSA
- 4. Incubate for 45 minutes at 4°C in dark
- 5. Wash cells 2 3 x with cold PBS; centrifugation at 200 x g for 5 minutes each
- 6. Resuspend cells and analyse directly by flow cytometry; for later analysis resuspend cells in 1% neutral buffered formalin, store in dark at 4°C and analyse within 24 hours



Indirect immunofluorescent staining of human dermal fibroblasts.

Indirect immunofluorescence with AS02 and antiprolylhydroxylase antibody on cultured human dermal fibroblasts; staining with AS02 followed by goat anti-mouse IgG DTAF conjugate (green fluorescence) and the antiprolylhydroxylase antibody with goat anti-mouse CY3 conjugate (yellow fluorescence). (magnification x 300)

AS02 homogenously stains the entire cell surface, including the numerous pseudopods. A clustering or intensification of staining at cell-cell contact sites was not demonstrated.

Confocal laser scanning microscopic analysis shows a sharp AS02 labeling of the cell membrane and a diffuse cytoplasmatic staining, suggesting that the antigen occurs predominantly at the cell surface, but also in the cytoplasm.

For in vitro research use only. Not for diagnostic or therapeutic use.



BIOZOL GmbH Leipziger Str.4 D-85386 Eching Germany

www.biozol.de info@biozol.de +49 (0)89 - 3799 666 6





Storage and Stability

The antibody is stable for 9 months when stored as undiluted liquid at 2-8°C.

References

- 1. Saalbach A, Aneregg U, Bruns M, Schnabel E, Herrmann K, Haustein UF. Novel fibroblast-specific monoclonal antibodies: properties and specificities. *J Invest Dermatol.* 106(6):1314-9, 1996.
- 2. Saalbach A, Aust G, Haustein UF, Herrmann K, Anderegg U. The fibroblast-specific MAb AS02: a novel tool for detection and elimination of human fibroblasts. *Cell Tissue Res* 290(3):593-9, 1997.
- 3. Saalbach A, Kraft R, Herrmann K, Haustein UF, Anderegg U. The monoclonal antibody AS02 recognizes a protein on human fibroblasts being highly homologous to Thy-1. *Arch Dermatol Res* 290(7):360-6, 1998.
- 4. Saalbach A, Wetzig T, Haustein UF, Anderegg U. Detection of human soluble Thy-1 in serum by ELISA. Fibroblasts and activated endothelial cells are a possible source of soluble Thy-1 in serum. *Cell Tissue Res* 298(2):307-15, 1999.
- 5. Saalbach A, Haustein UF, Anderegg U. A ligand of human thy-1 is localized on polymorphonuclear leukocytes and monocytes and mediates the binding to activated thy-1-positive microvascular endothelial cells and fibroblasts. *J Invest Dermatol* 115(5):882-8, 2000.
- 6. Hagood JS, Miller PJ, Lasky JA, Tousson A, Guo B, Fuller GM, McIntosh JC. Differential expression of platelet-derived growth factor-alpha receptor by Thy-1(-) and Thy-1(+) lung fibroblasts. *Am J Physiol* 277(1 Pt 1):L218-24, 1999.
- 7. Steiniger B, Barth P, Hellinger A. The perifollicular and marginal zones of the human splenic white pulp: do fibroblasts guide lymphocyte immigration? *Am J Pathol* 159(2):501-12, 2001.
- 8. Seemayer CA et al. Cartilage destruction mediated by synovial fibroblasts does not depend on proliferation in rheum atoid arthritis. *Am J Pathol* 162(5):1549-57, 2003.
- 9. Kollert, F et al. Soluble CD90 as a Potential Marker of Pulmonary. Involvement in Systemic Sclerosis. Arthritis Care & Research 65(2): 281–287, 2013.
- 10. Picke, A-K et al. Thy-1 (CD90) promotes bone formation and protects against obesity. Sci. Transl. Med. 10: 1-12, 2018
- 11. Saalbach, A. et al. Antifibrotic Soluble Thy-1 Correlates with Renal Dysfunction in Chronic Kidney Disease. *Int. J. Mol. Sci.* 24(3):1896m 2023, doi: 10.3390/ijms24031896.





